



Extensive cardinal parameter model to predict growth of pseudomonads in salt-reduced lightly preserved seafood

Martinez Rios, Veronica; Dalgaard, Paw

Publication date:
2017

Document Version
Publisher's PDF, also known as Version of record

[Link back to DTU Orbit](#)

Citation (APA):
Martinez Rios, V., & Dalgaard, P. (2017). *Extensive cardinal parameter model to predict growth of pseudomonads in salt-reduced lightly preserved seafood*. Abstract from 10th International Conference on Predictive Modelling in Food, Cordoba, Spain.

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

ABSTRACT

Reference:	84
Title:	Extensive cardinal parameter model to predict growth of pseudomonads in salt-reduced lightly preserved seafood
Author:	<u>Veronica Martinez-Rios</u> ¹ , Paw Dalgaard ¹
Institution/company:	1 National Food Institute (DTU Food) Technical University of Denmark Henrik Dams Allé, building 204 DK-2800 Kgs. Lyngby, Denmark
Topic:	PREDICTIVE MODELS FOR FOOD SAFETY AND QUALITY: DECONTAMINATION, FOOD FORMULATION,... ETC
Type of Communication:	ORAL
Keyword:	organic acids, product development
Introduction and Objectives:	<p>Interest in and demand for preserved seafood with reduced salt/sodium content is increasing. As a consequence of the reduced salt content potential growth of psychrotolerant pseudomonads to unacceptable high concentration where they cause product spoilage is an increasing challenge. Innovation is needed to reformulate these salt-reduced products and this must be done in such a way that other product characteristics compensate for less inhibiting effect due to salt. Numerous simple predictive models are available to predict growth of pseudomonads in foods at different temperatures. A few models include the effect of temperatures and salt. However, these simple secondary models do not include the effect of a broader range of product characteristics and therefore they cannot be used to predict how the inhibiting effect of salt can be replaced by changes in other environmental factors</p> <p>The objective was to develop an extensive predictive model that allows growth of psychrotolerant pseudomonads to be predicted in brined and marinated seafood with a range of different organic acids</p>
Materials and Methods:	<p>The new model was developed by expanding an existing cardinal parameter-type model for growth of pseudomonads in dairy products and including terms for temperature, pH, aw/NaCl, lactic- and sorbic acids (Martinez-Rios et al., Int. J. Food Microbiol. 216: 110-120, 2016). MIC-values for acetic-, benzoic- and citric acids were determined in broth and terms modelling their antimicrobial effect were added to the model. The new and expanded model included eight environmental factors and their interactive effects. The new model was evaluated under constant and dynamic temperature conditions using challenge tests and storage trials with a total of 78 growth curves in well characterized seafoods including lumpfish roe and brined shrimps</p>
Result:	<p>Average bias and accuracy factor values were 1.07 and 1.20 for 69 growth curves at constant temperatures. For nine growth curves at dynamic temperatures 77 % of the growth data (log cfu/g) was within the acceptable simulation zone of +/- 0.5 log cfu/g. The new model can be used to facilitate product reformulation as shown here for brined shrimps at 8°C, pH of 5.8 and water phase organic acid concentrations of 3000 ppm (citric), 1200 ppm (benzoic) and 500 ppm (sorbic). When the water phase salt concentration in this product is reduced from 3% to 1% growth of psychrotolerant pseudomonads change from none in 42 days to > 7 log cfu/g increase in 8 days. However, by including 500 ppm of acetic acid the reformulated prevent growth of psychrotolerant pseudomonads</p>
Conclusion:	<p>The high number of environmental factors included in the new model makes it flexible and suitable to support product development as the effect of substituting one combination of preservatives with another can be predicted</p>